Neuroscience Letters, 143 (1992) 275–278
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Aarnisalo, A., see Kivipelto, L., 190
Adachi, T., Baramidze, D.G. and Sato, A., Stimulation of the nucleus basalis of Meynert increases cortical cerebral blood flow without influencing diameter of the pial artery in rats, 173
Adamsson, M., see Wahlestedt, C., 123
Ally, A., see Park, G.A.S., 259
Alvarez, R.B., see Askanas, V., 96
Andersson, D., see Wahlestedt, C., 123
Askanas, V., Engel, W.K. and Alvarez, R.B., Strong immunoreactivity of β-amyloid precursor protein, including the β-amyloid protein sequence, at human neuromuscular junctions, 96
Avoli, M., see Hwa, G.G.C., 83

Baramidze, D.G., see Adachi, T., 173
Bayliss, D.A., see Berger, A.J., 164
Bazzani, C., see Guarini, S., 181
Belai, A., Schmidt, H.H.H.W., Hoyle, C.H.V., Hassall, C.J.S., Saffrey, M.J., Moss, J., Förstermann, U., Murad, F. and Burnstock, G., Colocalization of nitric oxide synthase and NADPH-diaphorase in the myenteric plexus of the rat gut, 60
Belai, A., see Hassall, C.J.S., 65
Belaidi, M., see Ben Attia, M., 74
Belfiore, P., Ponzio, F., Biagetti, R., Berettera, C., Magnani, M. and Pozzi, O., Oxiracetam prevents the hippocampal cholinergic hypofunction induced by the NMDA receptor blocker AP7, 127
Ben Attia, M., N'Gouemo, P., Belaidi, M., Rondouin, G. and Chicheportiche, R., Kindling and electrode effects on the benzodiazepine receptors density of olfactory bulb and hippocampus after olfactory bulb kindling, 74
Bennett, J.P., see Snow, H., 119
Berettera, C., see Belfiore, P., 127
Berg, P., see Heuser-Link, M., 31
Berger, A.J., Bayliss, D.A. and Viana, F., Modulation of neonatal rat hypoglossal motoneuron excitability by serotonin, 164
Bertolini, A., see Guarini, S., 181
Biagetti, R., see Belfiore, P., 127
Birbaumer, N., see Lutzenberger, W., 10
Blakemore, C., see Nicoll, A., 5
Block, F., see Schwarz, M., 105
Böttger, B., see Finger, T.E., 15
Bosler, O., Vuillon-Cacciuttolo, G. and Saïdi, H., Long-term serotonin reinnervation of the suprachiasmatic nucleus after 5,7-dihydroxytryptamine axotomy in the adult rat, 159
Bowen, D.M., see Stratmann, G.C., 43
Buchanan, J.T., see Meer, D.P., 1
Burnstock, G., see Belai, A., 60
Burnstock, G., see Hassall, C.J.S., 65
Butler, S.H., see Persson, S., 247

Card, J.P., see Nadelhaft, I., 271
Chan, J., see Lustig, H.S., 229
Chandler, H.C., King, V., Corwin, J.V. and Reep, R.L., Thalamocortical connections of rat posterior parietal cortex, 237
Chávez, M., see Fisher, G.H., 215
Chetyrbok, I.S., see Lyamin, O.I., 263
Chicheportiche, R., see Ben Attia, M., 74
Colombari, E., see De Almeida Colombari, D.S., 255
Conway, B.A., see Kiehn, O., 243
Corwin, J.V., see Chandler, H.C., 237
Costa, E., see Rothstein, J.D., 210
Cusano, G.P., see Fisher, G.H., 215

D'Aniello, A., see Fisher, G.H., 215
De Almeida Colombari, D.S., Colombari, E., Saad, W.A., De Arruda Camargo, L.A., Renzi, A., De Luca Jr., L.A. and Vanderlei Menani, J., Effect of furosemide treatment on the central and peripheral pressor responses to cholinergic and adrenergic agonists, angiotensin II, hypertonic solution and vasopressin, 255
De Arruda Camargo, L.A., see De Almeida Colombari, D.S., 255
De Luca Jr., L.A., see De Almeida Colombari, D.S., 255
Del Bianco, E., see Santicioli, P., 19
Deng, Q.-S. and Johanson, C.E., Cyclic AMP alteration of chloride transport into the choroid plexus-cerebrospinal fluid system, 146
Di Pasquale, E., Morin, D., Monteau, R. and Hilaire, G., Serotonergic modulation of the respiratory rhythm generator at birth: an in vitro study in the rat, 91
Dietrich, H., see Zernig, G., 169
Dirlich, G., see Heuser-Link, M., 31

Edvinsson, L., see Wahlestedt, C., 123
Edwards, R.J., see McKenzie, J.E., 23
Egger, C., see Kirchmair, R., 143
Elbert, T., see Lutzenberger, W., 10
Engel, W.K., see Askanas, V., 96
Eriksson, P.S., Nilsson, M., Wågberg, M., Rönnbäck, L. and Hansson, E., Volume regulation of single astroglial cells in primary culture, 195

Ferrari, W., see Guarini, S., 181
Finger, T.E. and Böttger, B., Expression of the dopaminergic phenotype in the olfactory bulb: neither calcitonin gene-related peptide nor olfactory input is necessary, 15
Fischer-Colbrie, R., see Kirchmair, R., 143
Fisher, G.H., D'Aniello, A., Vetere, A., Cusano, G.P., Chávez, M. and Petrucelli, L., Quantitfication of D-aspartate in normal and Alzheimer brains, 215
Flor, H., see Lutzenberger, W., 10
Förstermann, U., see Belai, A., 60
Förstermann, U., see Hassall, C.J.S., 65
Francis, P.T., see Stratmann, G.C., 43

Galey, D. and Jaffard, R., Post-training medial septal stimulation improves spatial information processing in BALB/c mice, 87
Gee, P., see Kirchmair, R., 143
Gentleman, S.M., see McKenzie, J.E., 23
Geppetti, P., see Santicioli, P., 19
Goto, S., Nagahiro, S., Ushio, Y. and Hirano, A., Calcineurin, a calcium/calmodulin-regulated protein phosphatase, in mammalian neuroendocrine cells and neoplasms, 51
Greenberg, D.A., see Lustig, H.S., 229
Guarini, S., Bazzani, C., Tagliavini, S., Bertolini, A. and Ferrari, W., Capsaicin prevents the adrenocorticotropin-induced improvement of

cardiovascular function and survival in hemorrhage-shocked rats, 181
Guidotti, A., see Rothstein, J.D., 210

Hansson, E., see Eriksson, P.S., 195
Hara, M., Inoue, M., Yasukura, T., Ohnishi, S., Mikami, Y. and Inagaki, C., Uneven distribution of intracellular Cl^- in rat hippocampal neurons, 135
Hassall, C.J.S., Saffrey, M.J., Belai, A., Hoyle, C.H.V., Moules, E.W., Moss, J., Schmidt, H.H.H.W., Murad, F., Förstermann, U. and Burnstock, G., Nitric oxide synthase immunoreactivity and NADPH-diaphorase activity in a subpopulation of intrinsic neurones of the guinea-pig heart, 65
Hassall, C.J.S., see Belai, A., 60
Hatanaka, H., see Kushima, Y., 110
Heuser-Link, M., Dirlich, G., Berg, P., Vogl, L. and Scherg, M., Eyeblinks evoke potentials in the occipital brain region, 31
Hilaire, G., see Di Pasquale, E., 91
Hirano, A., see Goto, S., 51
Hogue-Angeletti, R., see Kirchmair, R., 143
Hoyle, C.H.V., see Hassall, C.J.S., 65
Hoyle, C.H.V., see Belai, A., 60
Hultborn, H., see Kiehn, O., 243
Hu-Tsai, M., Winter, J. and Woolf, C.J., Regional differences in the distribution of capsaicin-sensitive target-identified adult rat dorsal root ganglion neurons, 251
Huynh, D.P., Lin, C.T. and Pulst, S.M., Expression of neurofibromin, the neurofibromatosis 1 gene product: studies in human neuroblastoma cells and rat brain, 233
Hwa, G.G.C. and Avoli, M., Excitatory synaptic transmission mediated by NMDA and non-NMDA receptors in the superficial/middle layers of the epileptogenic human neocortex maintained in vitro, 83

Ikeda, K., see Nishimura, M., 35
Inagaki, C., see Hara, M., 135
Inanami, O., Ohno, K. and Sato, A., Responses of regional cerebral blood flow to intravenous administration of thyrotropin releasing hormone in aged rats, 151
Inoue, M., see Hara, M., 135
Ishizuka, K. and Murakami, T., Convergence of excitatory inputs from the chorda tympani, glossopharyngeal, and vagus nerves onto inferior salivatory nucleus neurons in the cat, 155
Isomura, G., see Karasawa, N., 267

Jaffard, R., see Galey, D., 87
Johanson, C.E., see Deng, Q.-S., 146

Kaba, H., see Li, C.-S., 48
Karasawa, N., Isomura, G. and Nagatsu, I., Production of specific antibody against L-DOPA and its ultrastructual localization of immunoreactivity in the house-shrew (*Suncus murinus*) lateral habenular nucleus, 267
Kataoka, Y., see Motomura, M., 139
Keast, J.R., A possible neural source of nitric oxide in the rat penis, 69
Kerwin, J.M., Morris, C.M., Perry, R.H. and Perry, E.K., Hippocampal nerve growth factor receptor immunoreactivity in patients with Alzheimer's and Parkinson's disease, 101
Kiehn, O., Hultborn, H. and Conway, B.A., Spinal locomotor activity in acutely spinalized cats induced by intrathecal application of noradrenaline, 243
King, V., see Chandler, H.C., 237
Kirchmair, R., Egger, C., Gee, P., Hogue-Angeletti, R., Fischer-Colbrie, R., Laslop, A. and Winkler, H., Differential subcellular distribution of PC1, PC2 and furin in bovine adrenal medulla and secretion of PC1 and PC2 from this tissue, 143
Kivipelto, L., Aarnisalo, A. and Panula, P., Neuropeptide FF is colocalized with catecholamine-synthesizing enzymes in neurons of the nucleus of the solitary tract, 190
Kollack, S.S. and Newman, S.W., Mating behavior induces selective expression of Fos protein within the chemosensory pathways of the male Syrian hamster brain, 223
Kushima, Y. and Hatanaka, H., Interleukin-6 and leukemia inhibitory factor promote the survival of acetylcholinesterase-positive neurons in culture from embryonic rat spinal cord, 110

Laakso, M.-L., see Lindroos, O.F.C., 219
Larkman, A., see Nicoll, A., 5
Laslop, A., see Kirchmair, R., 143
Lees, G.J. and Leong, W., The non-NMDA glutamate antagonist NBQX blocks the local hippocampal toxicity of kainic acid, but not the diffuse extrahippocampal damage, 39
Leinonen, L.M., see Lindroos, O.F.C., 219
Leong, W., see Lees, G.J., 39
Leysen, J.E., see Pauwels, P.J., 27
Li, C.-S., Kaba, H., Saito, H. and Seto, K., Oestrogen infusions into the amygdala potentiate excitatory transmission from the accessory olfactory bulb to tuberoinfundibular arcuate neurones in the mouse, 48
Lin, C.T., see Huynh, D.P., 233
Lindroos, O.F.C., Leinonen, L.M. and Laakso, M.-L., Melatonin binding to the anteroventral and anterodorsal thalamic nuclei in the rat, 219
Lowrie, M.B., see Snow, H., 119
Lustig, H.S., von Brauchitsch, K.L., Chan, J. and Greenberg, D.A., A novel inhibitor of glutamate release reduces excitotoxic injury in vitro, 229
Lutzenberger, W., Birbaumer, N., Flor, H., Rockstroh, B. and Elbert, T., Dimensional analysis of the human EEG and intelligence, 10
Lyamin, O.I. and Chetyrbok, I.S., Unilateral EEG activation during sleep in the Cape fur seal, *Arctocephalus pusillus*, 263

Maggi, C.A., see Zernig, G., 169
Maggi, C.A., see Santicioli, P., 19
Magnani, M., see Belfiore, P., 127
Mailleux, P., Parmentier, M. and Vanderhaeghen, J.-J., Distribution of cannabinoid receptor messenger RNA in the human brain: an in situ hybridization histochemistry with oligonucleotides, 200
McKenzie, J.E., Gentleman, S.M., Royston, M.C., Edwards, R.J. and Roberts, G.W., Quantification of plaque types in sulci and gyri of the medial frontal lobe in patients with Alzheimer's disease., 23
Meer, D.P. and Buchanan, J.T., Apamin reduces the late afterhyperpolarization of lamprey spinal neurons, with little effect on fictive swimming, 1
Mikami, Y., see Hara, M., 135
Miselis, R.R., see Nadelhaft, I., 271
Monteau, R., see Di Pasquale, E., 91
Morin, D., see Di Pasquale, E., 91
Morris, C.M., see Kerwin, J.M., 101
Moss, J., see Belai, A., 60
Moss, J., see Hassall, C.J.S., 65
Motomura, M., Niwa, M., Kataoka, Y., Ootsuru, I., Yoshimura, T., Tsujihata, M. and Nagataki, S., In vitro quantitative autoradiography of [^{125}I]α-bungarotoxin binding at the motor end-plates of experimental autoimmune myasthenia gravis, 139
Moules, E.W., see Hassall, C.J.S., 65

Mulkey, R.M. and Zucker, R.S., Monensin can transport calcium across cell membranes in a sodium independent fashion in the crayfish *Procambarus clarkii, 115*
Murad, F., see Belai, A., 60
Murad, F., see Hassall, C.J.S., 65
Murakami, T., see Ishizuka, K., 155
Murtha, S.M., see Park, G.A.S., 259

N'Gouemo, P., see Ben Attia, M., 74
Nadelhaft, I., Vera, P.L., Card, J.P. and Miselis, R.R., Central nervous system neurons labelled following the injection of pseudorabies virus into the rat urinary bladder, 271
Nagahiro, S., see Goto, S., 51
Nagataki, S., see Motomura, M., 139
Nagatsu, I., see Karasawa, N., 267
Namba, Y., see Nishimura, M., 35
Newman, S.W., see Kollack, S.S., 223
Nicoll, A., Larkman, A. and Blakemore, C., EPSPs in rat neocortical pyramidal neurones in vitro are prolonged by NMDA receptor-mediated currents, 5
Nilsson, M., see Eriksson, P.S., 195
Nishi, S., see Yoshimura, M., 131
Nishimura, M., Namba, Y., Ikeda, K. and Oda, M., Glial fibrillary tangles with straight tubules in the brains of patients with progressive supranuclear palsy, 35
Niwa, M., see Motomura, M., 139
Nyberg, F., see Persson, S., 247

Oda, M., see Nishimura, M., 35
Ohnishi, S., see Hara, M., 135
Ohno, K., see Inanami, O., 151
Ootsuru, I., see Motomura, M., 139

Pai, K.S. and Ravindranath, V., Quisqualic acid-induced neurotoxicity is protected by NMDA and non-NMDA receptor antagonists, 177
Panula, P., see Kivipelto, L., 190
Pappas, B.A., see Park, G.A.S., 259
Park, G.A.S., Pappas, B.A., Murtha, S.M. and Ally, A., Enriched environment primes forebrain choline acetyltransferase activity to respond to learning experience, 259
Parmentier, M., see Mailleux, P., 200
Pauwels, P.J. and Leysen, J.E., Blockade of nitric oxide formation does not prevent glutamate-induced neurotoxicity in neuronal cultures from rat hippocampus, 27
Perry, E.K., see Kerwin, J.M., 101
Perry, R.H., see Kerwin, J.M., 101
Persson, S., Post, C., Weil-Fugazza, J., Butler, S.H. and Nyberg, F., Decreased cerebrospinal fluid neuropeptide-converting enzyme activity in monoarthritic rats, 247
Petrucelli, L., see Fisher, G.H., 215
Ponzio, F., see Belfiore, P., 127
Post, C., see Persson, S., 247
Pozzi, O., see Belfiore, P., 127
Priestley, J.V., see Rattray, M., 55
Pritz, M.B. and Stritzel, M.E., Interconnections between the pretectum and visual thalamus in reptiles, *Caiman crocodilus*, 205
Procter, A.W., see Stratmann, G.C., 43
Pulst, S.M., see Huynh, D.P., 233

Racagni, G., see Tromba, C., 185
Rattray, M., Savery, D., Wotherspoon, G., Priestley, J.V. and Smith, G.S., Two populations of cells that express preprocholecystokinin mRNA in ventral periaqueductal grey, 55
Ravindranath, V., see Pai, K.S., 177
Reep, R.L., see Chandler, H.C., 237
Reis, D.J., see Wahlestedt, C., 123
Renzi, A., see De Almeida Colombari, D.S., 255
Roberts, G.W., see McKenzie, J.E., 23
Rockstroh, B., see Lutzenberger, W., 10
Rondouin, G., see Ben Attia, M., 74
Rönnbäck, L., see Eriksson, P.S., 195
Rothstein, J.D., Guidotti A. and Costa, E., Release of endogenous benzodiazepine receptor ligands (endozepines) from cultured neurons, 210
Royston, M.C., see McKenzie, J.E., 23

Saad, W.A., see De Almeida Colombari, D.S., 255
Saffrey, M.J., see Hassall, C.J.S., 65
Saffrey, M.J., see Belai, A., 60
Saïdi, H., see Bosler, O., 159
Saito, H., see Li, C.-S., 48
Salvaggio, A., see Tromba, C., 185
Santicioli, P., Del Bianco, E., Geppetti, P. and Maggi, C.A., Release of calcitonin gene-related peptide-like (CGRP-LI) immunoreactivity from rat isolated soleus muscle by low pH, capsaicin and potassium, 19
Saria, A., see Zernig, G., 169
Sato, A., see Adachi, T., 173
Sato, A., see Inanami, O., 151
Savery, D., see Rattray, M., 55
Scherg, M., see Heuser-Link, M., 31
Schmidt, H.H.H.W., see Hassall, C.J.S., 65
Schmidt, H.H.H.W., see Belai, A., 60
Schwarz, M., Block, F. and Sontag, K.-H., *N*-Methyl-D-aspartate (NMDA)-mediated muscle relaxant action of memantine in rats, 105
Seto, K., see Li, C.-S., 48
Smith, G.S., see Rattray, M., 55
Snow, H., Lowrie, M.B. and Bennett, J.P., A postsynaptic GABA transporter in rat spinal motor neurones, 119
Sontag, K.-H., see Schwarz, M., 105
Stratmann, G.C., Webster, M.-T., Francis, P.T., Procter, A.W. and Bowen, D.M., Protease 'Clipsin' extract activity reflects demographic characteristics of human brain, 43
Stritzel, M.E., see Pritz, M.B., 205

Tagliavini, S., see Guarini, S., 181
Thomsen, R.H., see Wilson, D.F., 79
Tromba, C., Salvaggio, A., Racagni, G. and Volterra, A., Hypoglycemia-activated K+ channels in hippocampal neurons, 185
Tsujihata, M., see Motomura, M., 139

Ushio, Y., see Goto, S., 51

Vanderhaeghen, J.-J., see Mailleux, P., 200
Vanderlei Menani, J., see De Almeida Colombari, D.S., 255
Vera, P.L., see Nadelhaft, I., 271
Vetere, A., see Fisher, G.H., 215
Viana, F., see Berger, A.J., 164
Vogl, L., see Heuser-Link, M., 31
Volterra, A., see Tromba, C., 185
von Brauchitsch, K.L., see Lustig, H.S., 229
Vuillon-Cacciuttolo, G., see Bosler, O., 159

Wågberg, M., see Eriksson, P.S., 195
Wahlestedt, C., Reis, D.J., Yoo, H., Adamsson, M., Andersson, D. and Edvinsson, L., A novel inositol phosphate selectively inhibits vasoconstriction evoked by the sympathetic co-transmitters neuropeptide Y (NPY) and adenosine triphosphate (ATP), 123
Webster, M.-T., see Stratmann, G.C., 43
Weil-Fugazza, J., see Persson, S., 247
Wilson, D.F. and Thomsen, R.H., Effects of Hexamethonium on transmitter release from the rat phrenic nerve, 79
Winkler, H., see Kirchmair, R., 143
Winter, J., see Hu-Tsai, M., 251
Woolf, C.J., see Hu-Tsai, M., 251
Wotherspoon, G., see Rattray, M., 55
Yasukura, T., see Hara, M., 135
Yoo, H., see Wahlestedt, C., 123
Yoshimura, M. and Nishi, S., Excitatory amino acid receptors involved in primary afferent-evoked polysynaptic EPSPs of substantia gelatinosa neurons in the adult rat spinal cord slice, 131
Yoshimura, T., see Motomura, M., 139

Zernig, G., Dietrich, H., Maggi, C.A. and Saria, A., The substance P (NK1) receptor antagonist (±)-CP-96,345 causes sedation and motor impairment in Swiss albino mice in the black-and-white box behavioral paradigm, 169
Zucker, R.S., see Mulkey, R.M., 115